



Resistance of Salmonds Against Aeromonas Salmonicida

Host Genetics as a Main Player

Holten-Andersen, Lars; Dalsgaard, Inger; Buchmann, Kurt

Published in:
Book of abstracts

Publication date:
2011

[Link back to DTU Orbit](#)

Citation (APA):
Holten-Andersen, L., Dalsgaard, I., & Buchmann, K. (2011). Resistance of Salmonds Against Aeromonas Salmonicida: Host Genetics as a Main Player. In *Book of abstracts* DAFINET.
<http://www.dafinet.dk/DAFINET/Home.html>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**RESISTANCE OF SALMONIDS AGAINST *AEROMONAS SALMONICIDA*:
HOST GENETICS AS A MAIN PLAYER**

Lars Holten-Andersen^{1,2}, Inger Dalsgaard², Kurt Buchmann¹

¹ *Laboratory of Aquatic Pathobiology, Section of Biomedicine, Department of Veterinary Disease
Biology Faculty of Life Sciences, University of Copenhagen, Frederiksberg, Denmark*

² *National Veterinary Institute, Technical University of Denmark, Copenhagen V, Denmark*

Furunculosis, caused by *Aeromonas salmonicida*, continues to be a health problem for the growing salmonid aquaculture. Despite effective vaccination programs regular outbreaks occur at Danish trout farms calling for repeated antibiotic treatment. We hypothesized that a difference in natural susceptibility to this disease might exist between Baltic salmon and the widely used rainbow trout. Hence, a cohabitation challenge model was applied to investigate the relative susceptibility to infection with *Aeromonas salmonicida* in rainbow trout and Baltic salmon. The course of infection was monitored daily over a 30-day period post challenge and the results were summarized in mortality curves.

A. salmonicida was recovered from mortalities during the entire test period. At day 30 the survival was 6.2 % and 34.0 % for rainbow trout and Baltic salmon, respectively. Significant differences in susceptibility to *A. salmonicida* were demonstrated between the two salmonids and hazard ratio estimation between rainbow trout and Baltic salmon showed a 3.36 higher risk of dying from the infection in the former.

The finding that Baltic salmon carries a high level of natural resistance to furunculosis might raise new possibilities for salmonid aquaculture in terms of minimizing disease outbreaks and the use of antibiotics.

E-mail: lhoa@life.ku.dk